

I.B: Sommerfeld theory of metals

- So far we have mostly considered a single electron interacting with a simple periodic potential
- Considering the more realistic situation of interacting electrons and nuclei of the crystal will be postponed for now
- We will consider a simpler case: **Sommerfeld model**

* Assume we have some "conduction" electrons in a metal

- We discussed a while ago that some electrons are tightly-bound to atoms while others may participate in bonding / conduction ("valence")

* Assume that these electrons feel a constant potential throughout the crystal:

$$\left[\frac{\vec{p}^2}{2m} + E_c \right] \psi(\vec{r}) = E \psi(\vec{r})$$

- All of the variations in crystal potential smeared out into constant E_c .

• Eigenfunctions are: $\psi_{\vec{k}}(\vec{r}) = \frac{1}{\sqrt{V}} e^{i\vec{k} \cdot \vec{r}}$

• Eigenvalues are: $E(\vec{k}) = E_c + \frac{\hbar^2 k^2}{2m}$

↑
Set to zero for simplicity



* Consider now N free electrons in volume V . We often define electron density with dimensionless parameter r_s :

$$\frac{4}{3} \pi r_s^3 a_B^3 = \frac{1}{n} = \frac{V}{N}$$

Bohr radius ↑ density

* Ground state means accommodating N electrons into N lowest energy available states

- Each state with wavevector \vec{k} , energy $E(\vec{k})$ can hold 2 electrons (up and down spin)
- States fill to Fermi energy E_F , Fermi wavevector k_F



- Fermi surface in \vec{k} space given by $E(\vec{k}) < E_F$ separates occupied and unoccupied states
- For free electron case, Fermi surface is a sphere in reciprocal space of radius k_F
- k_F given by requirement that total number of states equals N :

$$N = \sum_{\vec{k}} 2 = \frac{V}{(2\pi)^3} \int_{|\vec{k}| < k_F} d^3k \cdot 2 = 2 \frac{V}{(2\pi)^3} \frac{4}{3} \pi k_F^3 = \frac{V}{3\pi^2} k_F^3$$

↑
for spin

$$\Rightarrow k_F^3 = 3\pi^2 \frac{N}{V} = 3\pi^2 n$$

- Using the definition of r_s :

$$k_F^3 = 3\pi^2 \left(\frac{3}{4} \frac{1}{\pi r_s^3 a_B^3} \right) = \frac{9}{4} \frac{\pi}{r_s^3 a_B^3}$$

$$\Rightarrow k_F = \left(\frac{9\pi}{4} \right)^{1/3} \frac{1}{r_s a_B}$$

* Taking $p = \hbar k$, we can define a "Fermi velocity":

$$v_F = \frac{\hbar}{m} k_F$$

* Fermi energy: $E_F = \frac{\hbar^2 k_F^2}{2m} = \frac{\hbar^2}{2m a_B^2} \left(\frac{9\pi}{4}\right)^{2/3} \frac{1}{r_s^2}$

* We can then write the total energy of the free-electron gas as:

$$E_0 = 2 \sum_{\vec{k} < k_F} \frac{\hbar^2 k^2}{2m} = 2 \frac{V}{(2\pi)^3} \int_0^{k_F} 4\pi k^2 \frac{\hbar^2 k^2}{2m} dk = 8\pi \frac{V}{(2\pi)^3} \frac{\hbar^2}{2m} \frac{k_F^5}{5}$$

integrate over Fermi sphere

• Since $N = \frac{V}{3\pi^2} k_F^3$, energy per electron is:

$$\frac{E_0}{N} = \frac{\frac{V}{\pi^2} \frac{\hbar^2}{2m} \frac{k_F^5}{5}}{\frac{V}{3\pi^2} k_F^3} = \frac{3}{5} \frac{\hbar^2}{2m} k_F^2 = \frac{3}{5} E_F$$

* We can compute the DOS from the dispersion relation:

$$E(\vec{k}) = \frac{\hbar^2 k^2}{2m}$$

$$\Rightarrow D(E) = 2 \frac{V}{(2\pi)^3} \int \delta\left(E - \frac{\hbar^2 k^2}{2m}\right) d^3k$$

Recall $\delta[f(x)] = \sum_{x_0 \text{ zeros of } f} \frac{\delta(x-x_0)}{|f'(x_0)|}$


only + root in integration range

$$= \frac{2V}{(2\pi)^3} \int_0^{k_F} 4\pi k^2 \delta\left(E - \frac{\hbar^2 k^2}{2m}\right) dk$$

since $k_0 = \pm \sqrt{\frac{2mE}{\hbar^2}}$, $\frac{1}{|f'(k_0)|} = \frac{m}{\hbar^2 |k_0|}$

$$D(E) = \frac{2V}{(2\pi)^3} \cdot 4\pi \left(\frac{2mE}{\hbar^2} \frac{m}{\hbar^2} \frac{\hbar}{\sqrt{2mE}} \right) = \frac{V (2m)^{3/2}}{2\pi^2 \hbar^3} E^{1/2}$$

* DOS for free - electron gas:



- What about finite T?


* For an assembly of identical Fermions in thermal equilibrium, probability that one-particle state at energy E is occupied is Fermi-Dirac distribution:

$$f(E) = \frac{1}{e^{(E-E_F)/k_B T} + 1}$$


(see G and P Sec. III. Appendix B for derivation)

• $\frac{\partial f}{\partial E} = -\frac{e^{(E-E_F)/k_B T}}{[e^{(E-E_F)/k_B T} + 1]^2} \frac{1}{k_B T}$ is also a useful function (more later)

For $T=0$:



For $T > 0$:



* How does E_F change w/ T ?

- To calculate that, we perform Sommerfeld expansion
- Consider integral with a generic (smooth) function $G(E)$:

$$\int_{-\infty}^{\infty} G(E) \left(-\frac{\partial f}{\partial E}\right) dE$$

- Expand $G(E)$ around E_F :

$$G(E) = G(E_F) + (E-E_F) \left. \frac{dG}{dE} \right|_{E=E_F} + \frac{1}{2} (E-E_F)^2 \left. \frac{d^2 G}{dE^2} \right|_{E=E_F} + \dots$$

$$\frac{d^2 G}{dE^2} \Big|_{E=E_F}$$

- So integral becomes:

$$\int_{-\infty}^{\infty} G(E) \left(-\frac{\partial F}{\partial E}\right) dE = G(E_F) + \frac{1}{2} G''(E_F) \int_{-\infty}^{\infty} (E-E_F)^2 \left(-\frac{\partial F}{\partial E}\right) dE + \dots$$

where we used the fact that $\int_{-\infty}^{\infty} \left(-\frac{\partial F}{\partial E}\right) dE = -[F(\infty) - F(-\infty)] = 1$

And that odd powers give zero contribution, since $-\frac{\partial F}{\partial E}$ is even around E_F , so $(E-E_F)^{2n+1} \left(-\frac{\partial F}{\partial E}\right)$ is an odd function around E_F , integral over all E vanishes.

- Coeff. of the G'' is:

$$\frac{1}{2} \int_{-\infty}^{\infty} (E-E_F)^2 \left(-\frac{\partial F}{\partial E}\right) dE = \frac{1}{2} \int_{-\infty}^{\infty} (E-E_F)^2 \frac{e^{(E-E_F)/k_B T}}{[e^{(E-E_F)/k_B T} + 1]^2} \frac{1}{k_B T} dE$$

Change variables $(E-E_F)/k_B T \rightarrow x$ $dx = \frac{dE}{k_B T}$

and use the fact that it is even function: $\frac{1}{2} \int_{-\infty}^{\infty} \rightarrow \int_0^{\infty}$
 $= k_B^2 T^2 \int_0^{\infty} x^2 \frac{e^x}{(e^x+1)^2} dx = k_B^2 T^2 \frac{\pi^2}{6}$ (see Gr and P sec III.2)

$$\text{So: } \int_{-\infty}^{\infty} G(E) \left(-\frac{\partial F}{\partial E}\right) dE = G(E_F) + G''(E_F) k_B^2 T^2 \frac{\pi^2}{6} + \dots$$

- In a similar way can get $G'''(E_F)$ term, etc.

- Integrate the LHS by parts: $\int u dv = uv - \int v du$

$$\int_{-\infty}^{\infty} G(E) \left(-\frac{\partial F}{\partial E}\right) dE = -G(E) F(E) \Big|_{-\infty}^{\infty} + \int_{-\infty}^{\infty} \frac{dG(E)}{dE} F(E) dE$$

Assume: $G(-\infty) = 0$
 $G(\infty)$ well behaved
 Then $G(E) F(E) \Big|_{-\infty}^{\infty} = 0$

• Therefore!

$$\int_{-\infty}^{\infty} \frac{dG(E)}{dE} f(E) dE = G(E_F) + \frac{\pi^2}{6} k_B T^2 G''(E_F) + \dots$$

• define $\Gamma(E)$ s.t. $G(E) = \int_{-\infty}^E \Gamma(E') dE'$

then!

$$\int_{-\infty}^{\infty} \Gamma(E) f(E) dE = \int_{-\infty}^{E_F} \Gamma(E) dE + \frac{\pi^2}{6} (k_B T)^2 \left. \frac{d\Gamma(E)}{dE} \right|_{E=E_F} + \mathcal{O}(T^4)$$

• known as Sommerfeld expansion

• Rapidly convergent. For example consider $G(E) \sim E^p$ i.e., polynomial type.

$$\Rightarrow \text{Then: } \int_{-\infty}^{\infty} \frac{dG(E)}{dE} f(E) dE = G(E_F) + \frac{\pi^2}{6} p(p-1) \left(\frac{k_B T}{E_F} \right)^2 G''(E_F) + \dots$$

so terms decrease like $\left(\frac{k_B T}{E_F} \right)^{2n}$

$\Rightarrow E_F$ for most metals is ~ 10 eV, $k_B T \sim 0.026$ eV

• Now lets use Sommerfeld expansion on DOS at finite T :

$$\int_{-\infty}^{\infty} D(E) f(E) dE = N = \int_{-\infty}^{E_F} D(E) dE + \frac{\pi^2}{6} k_B^2 T^2 \left. \frac{dD(E)}{dE} \right|_{E=E_F} + \mathcal{O}(T^4)$$

neglect T^4 and higher terms

• Differentiate both sides by T , using that

$$\frac{d}{dT} \int_{-\infty}^{E_F(T)} D(E) dE = \frac{dE_F}{dT} D(E_F) + \int_{-\infty}^{E_F(T)} \frac{dD(E)}{dT} dE = \frac{dE_F}{dT} D(E_F)$$

$$\hookrightarrow \frac{d}{db} \int_a^b f(x) dx = f(b)$$

So:

$$\frac{dN}{dT} = 0 = \frac{dE_F}{dT} D(E_F) + \frac{\pi^2}{3} k_B^2 T \left. \frac{dD(E)}{dE} \right|_{E=E_F} + \mathcal{O}(T^2)$$

neglect

• So:

$$D'(E_F) = \left. \frac{dD(E)}{dE} \right|_{E=E_F}$$

$$\frac{dE_F}{dT} = -\frac{\pi^2}{3} k_B^2 T \frac{D'(E_F)}{D(E_F)}$$

• For 3D free electron gas, $D(E) \propto E^{3/2}$ so

$$\frac{D'(E_F)}{D(E_F)} = \frac{1}{2} \frac{1}{\sqrt{E_F}} = \frac{1}{2E_F} \quad \text{and} \quad \frac{dE_F}{dT} = -\frac{\pi^2}{6} k_B^2 T \frac{1}{E_F}$$

$$E_F \frac{dE_F}{dT} = -\frac{\pi^2}{6} k_B^2 T$$

$$\frac{1}{2} \frac{d}{dT} (E_F^2) = -\frac{\pi^2}{6} k_B^2 T$$

Integrate both sides:

$$\frac{1}{2} \int_0^T \frac{d}{dT'} (E_F^2) dT' = -\frac{\pi^2}{6} k_B^2 \int_0^T T' dT'$$

$$\Rightarrow \frac{1}{2} [E_F^2(T) - E_F^2(0)] = -\frac{\pi^2}{12} k_B^2 T^2$$

$$\Rightarrow E_F(T) = \sqrt{E_F^2(0) - \frac{\pi^2}{6} k_B^2 T^2}$$

$$= E_F(0) \sqrt{1 - \frac{\pi^2}{6} \frac{k_B^2 T^2}{E_F^2(0)}}$$

Small! $\sqrt{1-x} = 1 - \frac{x}{2} - \dots$

$$E_F(T) \approx E_F(0) \left[1 - \frac{\pi^2}{12} \left(\frac{k_B T}{E_F(0)} \right)^2 \right]$$

↳ Slow decrease in E_F as a function of T

- We can analyze other properties of the free electron gas

- Consider the heat capacity: $C_V = \left. \frac{\delta Q}{dT} \right|_V$

↑ heat transferred from the environment (path/process function)
 ↑ T increase of system (state function)

* Extensive quantity (depends on system size)

* Consider the change in internal energy U :

$$dU = \delta Q + \delta L$$

↑ Work done on the system, $\delta L = -pdV$ if just mechanical work

Therefore, $C_V = \left. \frac{dU}{dT} \right|_V$ ← constant V

$$\begin{aligned}
 U(T) &= \int_{-\infty}^{\infty} E D(E) f(E) dE \\
 &= \int_{-\infty}^{E_F} E D(E) dE + \frac{\pi}{6} k_B^2 T^2 \left. \frac{d}{dE} [E D(E)] \right|_{E_F} + \mathcal{O}(T^4) \\
 &= \int_{-\infty}^{E_F} E D(E) dE + \frac{\pi}{6} k_B^2 T^2 \left[D(E_F) + E_F D'(E_F) \right]
 \end{aligned}$$

↑ energy of state times DOS

So using: $\frac{d}{dT} \int_{-\infty}^{E_F(T)} E D(E) dE = E_F D(E_F) \frac{dE_F}{dT}$,

$$\begin{aligned}
 C_V = \left. \frac{dU}{dT} \right|_V &= E_F D(E_F) \frac{dE_F}{dT} + \frac{\pi^2}{6} k_B^2 \left[2T D(E_F) + 2T E_F D'(E_F) \right. \\
 &\quad \left. + \cancel{\mathcal{O}(T^2)} \right]
 \end{aligned}$$

We know from before that: $\frac{dE_F}{dT} = -\frac{\pi^2}{3} k_B^2 T \frac{D'(E_F)}{D(E_F)}$

$$C_V = E_F D(E_F) \left[-\frac{\pi^2}{3} k_B^2 T \frac{D'(E_F)}{D(E_F)} \right] + \frac{\pi^2}{3} k_B^2 \left[T D(E_F) + T E_F D'(E_F) \right]$$

$$= \frac{\pi^2}{3} k_B^2 T D(E_F)$$

- Since E_F is weakly T dependent, we can write

$$C_V = \frac{\pi^2}{3} k_B^2 T D[E_F(0)]$$

Fermi energy at $T=0$

- Note, only electron contribution to C_V (will discuss lattice contribution later)

- For free electron gas, DOS $\propto D(E) = V \frac{(2m)^{3/2}}{2\pi^2 \hbar^3} E^{1/2}$

$$C_V = \frac{\pi^2}{3} k_B^2 T \frac{3}{2} \frac{N}{E_F} \leftarrow \text{number of } e^- \quad \text{and} \quad D(E_F) = \frac{3}{2} \frac{N}{E_F}$$

Define $T_F = \frac{E_F}{k_B} \Rightarrow C_V = \frac{\pi^2}{2} k_B \frac{T}{T_F}$

only electrons around E_F contribute to C_V

* If the heat is received as part of a reversible process in a closed system, 2nd law of thermodynamics states!

$$dS = \frac{\delta Q}{T} \quad \text{so} \quad C_V = T \left. \frac{dS}{dT} \right|_V$$

$$\bullet \text{ so: } \int_0^T \frac{dS}{dT'} dT' = \int_0^T \frac{C_V}{T'} dT' = \int_0^T \frac{\pi^2}{3} k_B^2 D[E_F(0)] dT'$$

$$= S(T) - S(0) = \frac{\pi^2}{3} k_B^2 T D[E_F(0)] = C_V$$

third law of thermodynamics

$\Rightarrow S = C_V$, entropy of electronic system is same as the heat capacity

Sommerfeld model of metals: What have we learned?

- Electrons are filled into energy states up to the Fermi energy.
- At finite temperature, the occupation of states is given by the Fermi distribution
- Temperature dependence of quantities like the Fermi energy, internal energy, etc. can be estimated by Sommerfeld expansion
 - * E_F is weakly T dependent since usually $E_F \gg k_B T$
- Heat capacity is the same as entropy in the electronic system.